FORM B.

Return showing the description and weight of all yarn spun in the Cotton Mills in the Mysore State, during the month ending the 31st January 1930.

Count of yarn	Weight in pounds of each count	Count of yarn	Weight in pounds of each count	Count of yarn	Weight in pounds of each count
4s	2,704	14½s	29,218	26s	26,840
5½s	1,993	16s	20,927	28s	1,08,761
6s	42,262	16½s	4,382	30s	61,057
6½s	6,131	17s	189	32s	6,587
8s	30,557	18s	21	40s	1,04,516
8½s	58,323	19s	1,55,806	60s	236
10s	24,274	20s	2,22,440		
10½s	20,974	20½s	5,284		
12s	31,668	21s	5,786		
12½s	29,383	22s	5,646		
13½s	19,588	22½s	3,246	Total ..	11,99,590
14s	43,183	24s	1,32,688		

BANGALORE,
Dated the 18th February 1930.

H. K. RAMIENGAR,
For *Director*.

OFFICE OF THE DIRECTOR OF HEALTH IN MYSORE, BANGALORE.

NOTE ON THE STATE OF PUBLIC HEALTH IN MYSORE, FOR THE WEEK ENDING SATURDAY, THE 15TH FEBRUARY 1930.

A total of 77 attacks with 63 deaths from plague was returned for the week, against 111 attacks with 70 deaths in the previous week. The Cities of Bangalore and Mysore returned 5 attacks with 4 deaths and one attack with one death respectively while the Kolar Gold Field area continued to remain free from the disease.

2. Comparative figures for the corresponding week for the State were 16 attacks with 12 deaths in 1928 and 122 attacks with 94 deaths in 1929.

3. One-hundred-and-fifty-four attacks with 31 deaths from small-pox were returned for 3 villages in Koppa Taluk, 2 villages in Tirthahalli Taluk, one village in each of the Mudgere, Malvalli, Tarikere, Honnali Taluks and Shimoga, Koppa, Haribar, Hosdurga, Chikmagalur, Kumsi and Birur Towns including Mysore and Bangalore Cities.

4. The State continued to remain free from other epidemic diseases, *viz.*, cholera, influenza and relapsing fever.

5. The total mortality for the Bangalore City amounted to 90, representing a calculated annual ratio of 39·94 per mille of population against the average ratio of 32·58 for 1928. Classified according to causes, the deaths numbered 4 under plague, 4 under malaria, 5 under broncho pneumonia, 6 under pneumonia, 7 under other fevers, 9 under phthisis, 6 under dysentery and diarrhœa and 49 under all other causes.

6. As compared with the City, the Civil and Military Station returned 93 deaths for the week ending 9th February 193), representing a calculated annual ratio of 42·99 per mille of population against 32·02 in the previous week. The causes of deaths were cholera 4, typhoid 1, malaria 1, other fevers 1, dysentery and diarrhoea 7, consumption 11, broncho pneumonia 9, pneumonia 11 and all other causes 53.

7. The total mortality for the Mysore City for the week was 53, representing a calculated annual ratio of 32 82 per mille of population against the average ratio of 33 51 for 1928. The causes of deaths were plague 1, malaria 2, pneumonia 4, typhoid 1, phthisis 5, dysentery and diarrhoea 7 and all other causes 35.

8. There were 39 deaths from all causes, reported from the Kolar Gold Field area equivalent to a calculated annual ratio of 23·12 per mille of population against the average ratio of 24·31 for 1928. The causes of deaths were broncho pneumonia 2, pneumonia 1, other fevers 1, phthisis 1, dysentery and diarrhœa 6 and all other causes 28.

J. V. KARVE,
Director of Health.